

Math 2B: midterm
Friday August 25 2017, 1:00 – 2:15pm

There are 9 exercises, worth a total of 100 points.
No electronic devices/books/notes allowed.
Provide computations and or explanations, unless stated otherwise.

Name:

Student ID:

Exercise 1 (5 points each) Integrate.

(a) $\int \frac{\sin(x)}{1+\cos^2(x)} \, dx$

(b) $\int x^2 \sin(2x) \, dx$

(c) $\int \tan^5(x) \sec^3(x) \, dx$

(d) $\int \frac{dx}{\sqrt{x^2+4}}$. You may use: $\int \sec(\theta) \, d\theta = \ln |\sec(\theta) + \tan(\theta)| + C$.

Exercise 2 (5 points each) Determine whether each improper integral below is convergent or divergent. Evaluate those that are convergent.

(a) $\int_1^5 (x-1)^{-4/3} \, dx$

(b) $\int_0^\infty x e^{-x} \, dx$

(c) $\int_3^\infty \frac{1}{x^2-x-2} \, dx.$

Exercise 3 (5 points each) Consider the function $f(x) = 2x$.

(a) Estimate, using Riemann sums, the area under $f(x)$ and above the x -axis for $x = 1$ to $x = 3$ using four rectangles and left-hand endpoints.

(b) Express $\int_1^3 f(x) \, dx$ as a limit of Riemann sums.

(c) Evaluate the limit from part b directly (do not use the fundamental theorem of calculus). One may use the formula: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$.

Exercise 4 (5 points) Find $g'(x)$ if $g(x) = \int_x^{\sin(x)} \frac{t^2+2t}{e^t+1} dt$.

Exercise 5

(a) (5 points) Find the area of the region enclosed by $y = x$ and $y = x^4$.

(b) (10 points) Find the volume of the solid generated by revolving the region described in part (a) about $y = -1$.


(c) (10 points) Find the volume of the solid whose base is the region described in part (a) and whose cross-sections perpendicular to the y -axis are squares with one side in the xy -plane.

Exercise 6 (5 points) Find the average value of the function $f(x) = \sin^3(x)$ on $[0, \pi]$.

Exercise 7 (5 points) Assume that f is a continuous function with $\int_1^4 f(x) \, dx = 42$. Compute $\int_1^2 xf(x^2) \, dx$.

Exercise 8 (5 points) A sail boat sails in a straight line so that its velocity at time t is $v(t) = t^3 - 1$. Find the *distance traveled* of the boat between $t = 0$ and $t = 2$.

Exercise 9: (5 points) Let $g(x) = \int_1^x f(t) \, dt$ where $f(t)$ is given by the following plot:



Fill in the blanks:

$$g(0) = \underline{\hspace{2cm}}$$

$$g(1) = \underline{\hspace{2cm}}$$

$$g(6) = \underline{\hspace{2cm}}$$

$$g'(2) = \underline{\hspace{2cm}}$$

$$g'(4) = \underline{\hspace{2cm}}.$$

Solutions:

1a: substitute $u = \cos(x)$. Then

$$\int \frac{\sin(x)}{1 + \cos^2(x)} dx = - \int \frac{1}{1 + u^2} du = -\tan^{-1}(u) + C = -\tan^{-1}(\cos(x)) + C.$$

1b: integration by parts (standard computation)

$$1/4 \cos(2x) - 1/2x^2 \cos(2x) + 1/2x \sin(2x) + C.$$

1c: rewrite and substitute $u = \sec(x)$:

$$\begin{aligned} \int \tan^5(x) \sec^3(x) dx &= \int (\sec^2(x) - 1)^2 \sec^2(x) \sec(x) \tan(x) dx \\ &= \int (u^2 - 1)^2 u^2 du = 1/7 u^7 - 2/5 u^5 + 1/3 u^3 + C \\ &= 1/7 \sec^7(x) - 2/5 \sec^5(x) + 1/3 \sec^3(x) + C \end{aligned}$$

1d: $x = 2 \tan(\theta)$ transforms the integral into $\int \sec(\theta) d\theta$. Then apply the given formula to obtain

$$\int \frac{dx}{\sqrt{x^2 + 4}} = \ln |\sqrt{x^2 + 4} + x| + C$$

2a: $-3(x-1)^{-1/3}|_1^5$, it diverges.

2b: $-xe^{-x} - e^{-x}|_0^\infty = 1$.

2c: Note: $1/(x^2 - x - 2) = 1/((x-2)(x+1)) = 1/3 \cdot 1/(x-2) - 1/3 \cdot 1/(x+1)$.

Hence integral is

$$\int_3^\infty 1/(x^2 - x - 2) dx = 1/3 \ln \left(\left| \frac{x-2}{x+1} \right| \right) \Big|_3^\infty = \ln(4)/3.$$

This integral diverges.

3a: $1/2 (f(1/2) + f(3/2) + f(2) + f(5/2)) = 1/2(2 + 3 + 4 + 5) = 7$.

3b: $\lim_{n \rightarrow \infty} 2/n \sum_{i=1}^n 2(1 + 2i/n)$ (also other options, this is right hand points).

3c: A simple computation shows that the limit becomes

$$\lim_{n \rightarrow \infty} 4 + 4(n+1)/n = 8.$$

4:

$$\cos(x) \frac{\sin(x)^2 + 2 \sin(x)}{e^{\sin(x)} + 1} - 3x^2 \frac{x^2 + 2x}{e^x + 1}.$$

5a: $\int_0^1 (x - x^4) dx = 3/10$.

5b: $\int_0^1 \pi((x+1)^2 - (x^4+1)^2) dx = 37/45\pi$.

5c: $\int_0^1 (y^{1/4} - y)^2 dy = 1/9$.

6:

$$1/\pi \int_0^\pi \sin^3(x) dx = 1/\pi \int_0^\pi (1 - \cos^2(x)) \sin(x) dx = \frac{1}{\pi} \cdot (-\cos(x) + 1/3 \cos^3(x)) \Big|_0^\pi = 4/3.$$

7: substitution:

$$\int_1^2 x f(x^2) du = \int_1^4 1/2 f(u) du$$

($u = x^2$) and hence the result is $1/2 \cdot 42 = 21$.

8: $\int_0^1 (1 - t^3) dt + \int_1^2 (t^3 - 1) dt = (t - t^4/4)|_0^1 + (t^4/4 - t)|_1^2 = 7/2$.

9: $g(0) = -2$, $g(1) = 0$, $g(6) = 1$, $g'(2) = 4$ and $g'(4) = -1$.