

# Bayesian Networks

Read R&N Ch. 13.6, 14.1-14.5

# You will be expected to know

- Basic concepts and vocabulary of Bayesian networks.
  - Nodes represent random variables.
  - Directed arcs represent (informally) direct influences.
  - Conditional probability tables,  $P(X_i | \text{Parents}(X_i))$ .
- Given a Bayesian network:
  - Write down the full joint distribution it represents.
- Given a full joint distribution in factored form:
  - Draw the Bayesian network that represents it.
- Given a variable ordering and some background assertions of conditional independence among the variables:
  - Write down the factored form of the full joint distribution, as simplified by the conditional independence assertions.
- Use the network to find answers to probability questions about it.

# Why Bayesian Networks?


---

- Probabilistic Reasoning
  - Knowledge Base : Joint distribution over all random variables
  - Reasoning: Compute probability of states of the world
    - Find the most probable assignments
    - Compute marginal / conditional probability
  
- Why Bayesian Net?
  - Manipulating full joint distribution is very hard!
  - Exploit conditional independence properties
  - Bayesian Network usually more compact & feasible
    - Probabilistic Graphical Models
    - Tool for Reasoning, Computation
    - Probabilistic Reasoning based on the Graph

## Conditional independence


---

- Recall: chain rule of probability
  - $p(x,y,z) = p(x) p(y|x) p(z|x,y)$
- *Some* of these models are conditionally independent
  - e.g.,  $p(x,y,z) = p(x) p(y|x) p(z|x)$
- *Some* models may have even *more* independence
  - E.g.,  $p(x,y,z) = p(x) p(y) p(z)$
- The more independence and conditional independence, the more compactly we can represent and reason over the joint probability distribution.




# Bayesian networks


- Directed graphical model
- Nodes associated with variables
- “Draw” independence in conditional probability expansion
  - Parents in graph are the RHS of conditional
- Ex:  $p(x, y, z) = p(x) p(y | x) p(z | y)$



- Ex:  $p(a, b, c, d) = p(a) p(b | a) p(c | a, b) p(d | b)$




Graph must be **acyclic**



Corresponds to an order over the variables (chain rule)

# Bayesian Network

- Specifies a joint distribution in a structured form:



- Dependence/independence shown by a directed graph:
  - Node = random variable
  - Directed Edge = conditional dependence
  - Absence of Edge = conditional independence

A	B	P(C)
t	t	0.2
t	f	0.4
f	t	0.3
f	f	0.3

- Allows concise view of joint distribution relationships:
  - Graph nodes and edges show conditional relationships between variables.
  - Tables provide probability data.

- **Tables are concise!!**

- $P(\neg A)$  is not shown since it can be inferred as  $(1 - P(A))$ , etc.

# Bayesian Networks

---

- Structure of the graph  $\Leftrightarrow$  Conditional independence relations

In general,

$$p(X_1, X_2, \dots, X_N) = \prod p(X_i \mid \text{parents}(X_i))$$


The full joint distribution

The graph-structured approximation

- Requires that graph is acyclic (no directed cycles)
- 2 components to a Bayesian network
  - The graph structure (conditional independence assumptions)
  - The numerical probabilities (for each variable given its parents)
- Also known as belief networks, graphical models, causal networks
- Parents in the graph  $\Leftrightarrow$  conditioning variables (RHS) in the formula

# Examples of 3-way Bayesian Networks

A, B, and C are independent.




**Marginal Independence:**  
 $p(A,B,C) = p(A) p(B) p(C)$

Parents in the graph  $\Leftrightarrow$   
conditioning variables (RHS)

# Examples of 3-way Bayesian Networks

A and B directly influence C.



Parents in the graph  $\Leftrightarrow$   
conditioning variables (RHS)

**Independent Causes:**

$$p(A,B,C) = p(C|A,B)p(A)p(B)$$


**“Explaining away” effect:**

**Given C, observing A makes B less likely  
e.g., earthquake/burglary/alarm example**

**A and B are (marginally) independent  
but become dependent once C is known**

# Examples of 3-way Bayesian Networks

A directly influences B;  
B directly influences C;  
but A influences C only  
indirectly through B.



Parents in the graph  $\Leftrightarrow$   
conditioning variables (RHS)


Markov dependence:  
 $p(A,B,C) = p(C|B) p(B|A)p(A)$

# Burglar Alarm Example

- Consider the following 5 binary variables:
  - B = a burglary occurs at your house
  - E = an earthquake occurs at your house
  - A = the alarm goes off
  - J = John calls to report the alarm
  - M = Mary calls to report the alarm
- What is  $P(B \mid M, J)$  ? (for example)
- We can use the full joint distribution to answer this question
  - Requires  $2^5 = 32$  probabilities
  - Can we use prior domain knowledge to come up with a Bayesian network that requires fewer probabilities?

# The Causal Bayesian Network

Generally, order variables so that resulting graph reflects assumed causal relationships.



Only requires 10 probabilities!

# Constructing a Bayesian Network: Step 1

- Order the variables in terms of influence (may be a partial order)

e.g.,  $\{E, B\} \rightarrow \{A\} \rightarrow \{J, M\}$

Generally, order variables to reflect the assumed causal relationships.

- Now, apply the chain rule, and simplify based on assumptions
- $P(J, M, A, E, B) = P(J, M \mid A, E, B) P(A \mid E, B) P(E, B)$

$$\approx P(J, M \mid A) \quad P(A \mid E, B) P(E) P(B)$$


$$\approx P(J \mid A) P(M \mid A) P(A \mid E, B) P(E) P(B)$$

These conditional independence assumptions are reflected in the graph structure of the Bayesian network

# Constructing this Bayesian Network: Step 2


- $P(J, M, A, E, B) = P(J | A) P(M | A) P(A | E, B) P(E) P(B)$

**Parents in the graph  $\Leftrightarrow$  conditioning variables (RHS)**



- There are 3 conditional probability tables (CPDs) to be determined:  
 $P(J | A)$ ,  $P(M | A)$ ,  $P(A | E, B)$ 
  - Requiring  $2 + 2 + 4 = 8$  probabilities
- And 2 marginal probabilities  $P(E)$ ,  $P(B)$  -> 2 more probabilities
- Where do these probabilities come from?
  - Expert knowledge
  - From data (relative frequency estimates)
  - Or a combination of both - see discussion in Section 20.1 and 20.2 (optional)

# The Resulting Bayesian Network




$$P(J, M, A, E, B) = P(J | A) P(M | A) P(A | E, B) P(E) P(B)$$

Generally, order variables so that resulting graph reflects assumed causal relationships.

# The Bayesian Network From a Different Variable Ordering

---




Parents in the graph  $\Leftrightarrow$  conditioning variables (RHS)

$$P(J, M, A, E, B) = P(E | A, B) P(B | A) P(A | M, J) P(J | M) P(M)$$

Generally, order variables so that resulting graph reflects assumed causal relationships.

# The Bayesian Network From a Different Variable Ordering

---




Parents in the graph  $\Leftrightarrow$  conditioning variables (RHS)

$$P(J, M, A, E, B) = P(A \mid B, E, M, J) P(B \mid E, M, J) P(E \mid M, J) P(J \mid M) P(M)$$

Generally, order variables to reflect the assumed causal relationships.

# Number of Probabilities Needed (1)

- Joint distribution



Full joint distribution:  
 $2^5 = 32$  probabilities

Structured distribution:  
 specify 10 parameters

E	B	A	J	M	P(...)
0	0	0	0	0	.93674
0	0	0	0	1	.00133
0	0	0	1	0	.00005
0	0	0	1	1	.00000
0	0	1	0	0	.00003
0	0	1	0	1	.00002
0	0	1	1	0	.00003
0	0	1	1	1	.00000
0	1	0	0	0	.04930
0	1	0	0	1	.00007
0	1	0	1	0	.00000
0	1	0	1	1	.00000
0	1	1	0	0	.00027
0	1	1	0	1	.00016
0	1	1	1	0	.00025
0	1	1	1	1	.00000

E	B	A	J	M	P(...)
1	0	0	0	0	.00946
1	0	0	0	1	.00001
1	0	0	1	0	.00000
1	0	0	1	1	.00000
1	0	1	0	0	.00007
1	0	1	0	1	.00004
1	0	1	1	0	.00007
1	0	1	1	1	.00000
1	1	0	0	0	.00050
1	1	0	0	1	.00000
1	1	0	1	0	.00000
1	1	0	1	1	.00000
1	1	1	0	0	.00063
1	1	1	0	1	.00037
1	1	1	1	0	.00059
1	1	1	1	1	.00000

## Number of Probabilities Needed (2)


- Consider  $n$  binary variables
- Unconstrained joint distribution requires  $O(2^n)$  probabilities
- If we have a Bayesian network, with a maximum of  $k$  parents for any node, then we need  $O(n 2^k)$  probabilities
- Example
  - Full unconstrained joint distribution
    - $n = 30, k = 4$ : need  $10^9$  probabilities for full joint distribution
  - Bayesian network
    - $n = 30, k = 4$ : need 480 probabilities

# Example of Answering a Simple Query

- What is  $P(\neg j, m, a, \neg e, b) = P(J = \text{false} \wedge M = \text{true} \wedge A = \text{true} \wedge E = \text{false} \wedge B = \text{true})$

$P(J, M, A, E, B) \approx P(J | A) P(M | A) P(A | E, B) P(E) P(B)$  ; by conditional independence


$$P(\neg j, m, a, \neg e, b) \approx P(\neg j | a) P(m | a) P(a | \neg e, b) P(\neg e) P(b) \\ \approx 0.10 \times 0.70 \times 0.94 \times 0.998 \times 0.001$$



# Hospital Alarm network

[Beinlich et al., 1989]


The “alarm” network: 37 variables, 509 parameters (rather than  $2^{37} = 10^{11}$  !)



## Reasoning in Bayesian networks

---

- Suppose we observe J
  - Observing J makes A more likely
  - A being more likely makes B more likely
- Suppose we observe A
  - Makes M more likely
- Observe A and J?
  - J doesn't add any more information about M
  - Observing A makes J, M independent
- How can we read independence directly from the graph?




## Reasoning in Bayesian networks

---

- How are J,M related given A?

- $P(M) = 0.0117$
- $P(M|A) = 0.7$
- $P(M|A,J) = 0.7$
- Conditionally independent

*(we actually know this by construction!)*




- Proof:

$$\begin{aligned} p(J, M|a) &\propto \sum_{e,b} p(e) p(b) p(a|e, b) p(J|a) p(M|a) \\ &= \left( \sum_{e,b} p(e, b, a) \right) p(J|a) p(M|a) \\ &= p(a) p(J|a) p(M|a) \\ &= c_a f_a(J) g_a(M) \end{aligned}$$

## Reasoning in Bayesian networks

---

- How are J,B related given A?
  - $P(B) = 0.001$
  - $P(B|A) = 0.3735$
  - $P(B|A,J) = 0.3735$
  - Conditionally independent




- Proof:

$$\begin{aligned} p(J, B|a) &\propto \sum_{e,m} p(e) p(B) p(a|e, B) p(J|a) p(m|a) \\ &= \left( \sum_e p(e, B, a) \right) p(J|a) \left( \sum_m p(m|a) \right) \\ &= p(B, a) p(J|a) \\ &= f_a(B) g_a(J) \end{aligned}$$

## Reasoning in Bayesian networks

---

- How are E,B related?
  - $P(B) = 0.001$
  - $P(B|E) = 0.001$
  - (Marginally) independent
- What about given A?
  - $P(B|A) = 0.3735$
  - $P(B|A,E) = 0.0032$
  - Not conditionally independent!
  - The “causes” of A become coupled by observing its value
  - Sometimes called “explaining away”



# Inference in Bayesian Networks

- $\mathbf{X} = \{ X1, X2, \dots, Xk \}$  = **query variables** of interest
  - $\mathbf{E} = \{ E1, \dots, El \}$  = **evidence variables** that are observed
    - ( $\mathbf{e}$ , an **event**)
  - $\mathbf{Y} = \{ Y1, \dots, Ym \}$  = **hidden variables** (nonevidence, nonquery)
- 
- **What is the posterior distribution of  $\mathbf{X}$ , given  $\mathbf{E}$ ?**
  - $P(\mathbf{X} | \mathbf{e}) = \alpha \sum_y P(\mathbf{X}, \mathbf{y}, \mathbf{e})$
- 
- **What is the most likely assignment of values to  $\mathbf{X}$ , given  $\mathbf{E}$ ?**
  - $\operatorname{argmax}_x P(\mathbf{x} | \mathbf{e}) = \operatorname{argmax}_x \sum_y P(\mathbf{x}, \mathbf{y}, \mathbf{e})$

# Inference in Bayesian Networks

## Simple Example

$P(A)$

.05

Disease1

$P(B)$

.02

Disease2



}

**Query Variables A, B**

}

**Hidden Variable C**

}

**Evidence Variable D**

A B  $P(C|A,B)$

t t .95

t f .90

f t .90

f f .005

C  $P(D|C)$

t .95

f .002

TempReg


Fever

**Note: Not an anatomically correct model of how diseases cause fever!**

Suppose that two different diseases influence some imaginary internal body temperature regulator, which in turn influences whether fever is present.

# Inference in Bayesian Networks

## Simple Example



What is the posterior conditional distribution of our query variables, given that fever was observed?

$$\begin{aligned}
 P(A,B|d) &= \alpha \sum_c P(A,B,c,d) \\
 &= \alpha \sum_c P(A)P(B)P(c|A,B)P(d|c) \\
 &= \alpha P(A)P(B) \sum_c P(c|A,B)P(d|c)
 \end{aligned}$$

$$\begin{aligned}
 P(a,b|d) &= \alpha P(a)P(b) \sum_c P(c|a,b)P(d|c) = \alpha P(a)P(b) \{ P(c|a,b)P(d|c) + P(\neg c|a,b)P(d|\neg c) \} \\
 &= \alpha .05 \times .02 \times \{ .95 \times .95 + .05 \times .002 \} \approx \alpha .000903 \approx .014
 \end{aligned}$$

$$\begin{aligned}
 P(\neg a,b|d) &= \alpha P(\neg a)P(b) \sum_c P(c|\neg a,b)P(d|c) = \alpha P(\neg a)P(b) \{ P(c|\neg a,b)P(d|c) + P(\neg c|\neg a,b)P(d|\neg c) \} \\
 &= \alpha .95 \times .02 \times \{ .90 \times .95 + .10 \times .002 \} \approx \alpha .0162 \approx .248
 \end{aligned}$$


$$\begin{aligned}
 P(a,\neg b|d) &= \alpha P(a)P(\neg b) \sum_c P(c|a,\neg b)P(d|c) = \alpha P(a)P(\neg b) \{ P(c|a,\neg b)P(d|c) + P(\neg c|a,\neg b)P(d|\neg c) \} \\
 &= \alpha .05 \times .98 \times \{ .90 \times .95 + .10 \times .002 \} \approx \alpha .0419 \approx .642
 \end{aligned}$$

$$\begin{aligned}
 P(\neg a,\neg b|d) &= \alpha P(\neg a)P(\neg b) \sum_c P(c|\neg a,\neg b)P(d|c) = \alpha P(\neg a)P(\neg b) \{ P(c|\neg a,\neg b)P(d|c) + P(\neg c|\neg a,\neg b)P(d|\neg c) \} \\
 &= \alpha .95 \times .98 \times \{ .005 \times .95 + .995 \times .002 \} \approx \alpha .00627 \approx .096
 \end{aligned}$$

$$\alpha \approx 1 / (.000903 + .0162 + .0419 + .00627) \approx 1 / .06527 \approx 15.32$$

# Inference in Bayesian Networks

## Simple Example



What is the most likely posterior conditional assignment of values to our query variables, given that fever was observed?

$$\begin{aligned}
 & \operatorname{argmax}_{\{a,b\}} P(a, b | d) \\
 &= \operatorname{argmax}_{\{a,b\}} \sum_c P(a, b, c, d) \\
 &= \{a, \neg b\}
 \end{aligned}$$

$$\begin{aligned}
 P(a, b | d) &= \alpha P(a)P(b) \sum_c P(c|a, b)P(d|c) = \alpha P(a)P(b) \{ P(c|a, b)P(d|c) + P(\neg c|a, b)P(d|\neg c) \} \\
 &= \alpha .05 \times .02 \times \{ .95 \times .95 + .05 \times .002 \} \approx \alpha .000903 \approx .014
 \end{aligned}$$

$$\begin{aligned}
 P(\neg a, b | d) &= \alpha P(\neg a)P(b) \sum_c P(c|\neg a, b)P(d|c) = \alpha P(\neg a)P(b) \{ P(c|\neg a, b)P(d|c) + P(\neg c|\neg a, b)P(d|\neg c) \} \\
 &= \alpha .95 \times .02 \times \{ .90 \times .95 + .10 \times .002 \} \approx \alpha .0162 \approx .248
 \end{aligned}$$


$$\begin{aligned}
 P(a, \neg b | d) &= \alpha P(a)P(\neg b) \sum_c P(c|a, \neg b)P(d|c) = \alpha P(a)P(\neg b) \{ P(c|a, \neg b)P(d|c) + P(\neg c|a, \neg b)P(d|\neg c) \} \\
 &= \alpha .05 \times .98 \times \{ .90 \times .95 + .10 \times .002 \} \approx \alpha .0419 \approx .642
 \end{aligned}$$

$$\begin{aligned}
 P(\neg a, \neg b | d) &= \alpha P(\neg a)P(\neg b) \sum_c P(c|\neg a, \neg b)P(d|c) = \alpha P(\neg a)P(\neg b) \{ P(c|\neg a, \neg b)P(d|c) + P(\neg c|\neg a, \neg b)P(d|\neg c) \} \\
 &= \alpha .95 \times .98 \times \{ .005 \times .95 + .995 \times .002 \} \approx \alpha .00627 \approx .096
 \end{aligned}$$

$$\alpha \approx 1 / (.000903 + .0162 + .0419 + .00627) \approx 1 / .06527 \approx 15.32$$

# Inference in Bayesian Networks

## Simple Example



What is the posterior conditional distribution of A, given that fever was observed? (I.e., temporarily make B into a hidden variable.)

We can use  $P(A,B|d)$  from above.

$$P(A|d) = \alpha \sum_b P(A,b|d)$$

$$\begin{aligned}
 P(a|d) &= \sum_b P(a,b|d) = P(a,b|d) + P(a,\neg b|d) \\
 &= (.014 + .642) \approx .656
 \end{aligned}$$


$$\begin{aligned}
 P(\neg a|d) &= \sum_b P(\neg a,b|d) = P(\neg a,b|d) + P(\neg a,\neg b|d) \\
 &= (.248 + .096) \approx .344
 \end{aligned}$$

This is a marginalization, so we expect from theory that  $\alpha = 1$ ; but check for round-off error.

A	B	$P(A,B d)$ from above
t	t	$\approx .014$
f	t	$\approx .248$
t	f	$\approx .642$
f	f	$\approx .096$

# Inference in Bayesian Networks

## Simple Example



What is the posterior conditional distribution of A, given that fever was observed, and that further lab tests definitely rule out Disease2? (I.e., temporarily make B into an evidence variable with B = false.)

$$P(A|\neg b,d) = \alpha P(A,\neg b|d)$$

$$P(a|\neg b,d) = \alpha P(a,\neg b|d) \approx \alpha .642 \approx .870$$

$$P(\neg a|\neg b,d) = \alpha P(\neg a,\neg b|d) \approx \alpha .096 \approx .130$$

$$\alpha \approx 1 / (.642 + .096) \approx 1 / .738 \approx 1.355$$

A	B	$P(A,B d)$ from above
t	t	$\approx .014$
f	t	$\approx .248$
t	f	$\approx .642$
f	f	$\approx .096$

# General Strategy for inference

---

- Want to compute  $P(q | e)$

Step 1:

$$P(q | e) = P(q,e)/P(e) = \alpha P(q,e), \quad \text{since } P(e) \text{ is constant wrt } Q$$

Step 2:

$$P(q,e) = \sum_{a..z} P(q, e, a, b, \dots, z), \quad \text{by the law of total probability}$$

Step 3:

$$\sum_{a..z} P(q, e, a, b, \dots, z) = \sum_{a..z} \prod_i P(\text{variable } i | \text{parents } i)$$

(using Bayesian network factoring)

Step 4:

Distribute summations across product terms for efficient computation

Section 14.4 discusses exact inference in Bayesian Networks. The complexity depends strongly on the network structure. The general case is intractable, but there are things you can do. Section 14.5 discusses approximation by sampling.


# Given a graph, can we “read off” conditional independencies?

## The “Markov Blanket” of $X$ (the gray area in the figure)

$X$  is conditionally independent of everything else, GIVEN the values of:

- \*  $X$ 's parents
- \*  $X$ 's children
- \*  $X$ 's children's parents


$X$  is conditionally independent of its non-descendants, GIVEN the values of its parents.




## D-Separation

---


- Prove sets  $X, Y$  independent given  $Z$ ?
- Check all *undirected* paths from  $X$  to  $Y$
- A path is “inactive” if it passes through:
  - (1) A “chain” with an observed variable



- (2) A “split” with an observed variable




- (3) A “vee” with **only unobserved** variables below it



- If all paths are inactive, conditionally independent!

# Naïve Bayes Model



$$P(C | X_1, \dots, X_n) = \alpha \prod P(X_i | C) P(C)$$

Features  $X$  are conditionally independent given the class variable  $C$

Widely used in machine learning

e.g., spam email classification:  $X$ 's = counts of words in emails

Probabilities  $P(C)$  and  $P(X_i | C)$  can easily be estimated from labeled data

# Naïve Bayes Model (2)

$$P(C | X_1, \dots, X_n) = \alpha \prod P(X_i | C) P(C)$$

Probabilities  $P(C)$  and  $P(X_i | C)$  can easily be estimated from labeled data

$$P(C = c_j) \approx \#(\text{Examples with class label } c_j) / \#(\text{Examples})$$

$$P(X_i = x_{ik} | C = c_j) \\ \approx \#(\text{Examples with } X_i \text{ value } x_{ik} \text{ and class label } c_j) \\ / \#(\text{Examples with class label } c_j)$$

Usually easiest to work with logs


$$\log [ P(C | X_1, \dots, X_n) ] \\ = \log \alpha + \sum [ \log P(X_i | C) + \log P(C) ]$$

**DANGER:** Suppose ZERO examples with  $X_i$  value  $x_{ik}$  and class label  $c_j$  ?  
An unseen example with  $X_i$  value  $x_{ik}$  will NEVER predict class label  $c_j$  !

Practical solutions: Pseudocounts, e.g., add 1 to every  $\#()$  , etc.

Theoretical solutions: Bayesian inference, beta distribution, etc.

# Hidden Markov Model (HMM)



Two key assumptions:

1. hidden state sequence is Markov
2. observation  $Y_t$  is CI of all other variables given  $S_t$

Widely used in speech recognition, protein sequence models


Since this is a Bayesian network polytree, inference is linear in  $n$

## Example: Car diagnosis

Initial evidence: car won't start

Testable variables (green), "broken, so fix it" variables (orange)

Hidden variables (gray) ensure sparse structure, reduce parameters



## Compact conditional distributions contd.

Noisy-OR distributions model multiple noninteracting causes

- 1) Parents  $U_1 \dots U_k$  include all causes (can add leak node)
- 2) Independent failure probability  $q_i$  for each cause alone

$$\Rightarrow P(X|U_1 \dots U_j, \neg U_{j+1} \dots \neg U_k) = 1 - \prod_{i=1}^j q_i$$


<i>Cold</i>	<i>Flu</i>	<i>Malaria</i>	$P(\text{Fever})$	$P(\neg \text{Fever})$
F	F	F	<b>0.0</b>	1.0
F	F	T	0.9	<b>0.1</b>
F	T	F	0.8	<b>0.2</b>
F	T	T	0.98	$0.02 = 0.2 \times 0.1$
T	F	F	0.4	<b>0.6</b>
T	F	T	0.94	$0.06 = 0.6 \times 0.1$
T	T	F	0.88	$0.12 = 0.6 \times 0.2$
T	T	T	0.988	$0.012 = 0.6 \times 0.2 \times 0.1$

Number of parameters **linear** in number of parents

# Examples of “real world” Bayesian Networks

## Genetic linkage analysis


---



- 6 individuals
- Haplotype: {2, 3}
- Genotype: {6}
- Unknown

# Examples of “real world” Bayesian Networks

## Pedigree model: 6 people, 3 markers



# Summary

- Bayesian networks represent a joint distribution using a graph
- The graph encodes a set of conditional independence assumptions
- Answering queries (or inference or reasoning) in a Bayesian network amounts to computation of appropriate conditional probabilities
- Probabilistic inference is intractable in the general case
  - Can be done in linear time for certain classes of Bayesian networks (polytrees: at most one directed path between any two nodes)
  - Usually faster and easier than manipulating the full joint distribution